# DEPARTMENT of ENVIRONMENTAL SERVICES Water Supply & Pollution Control Division - Biology Bureau

#### LAKE TROPHIC DATA

#### MORPHOMETRIC:

Lake: COFFIN POND	Lake Area (ha):	7.08
Town: SUGAR HILL	Maximum depth (m):	1.6
County: Grafton	Mean depth (m):	0.7
River Basin: Connecticut	Volume (m³):	47000
Latitude: 44°14'31" N	Relative depth:	0.6
Longitude: 71°45'53" W	Shore configuration:	2.39
Elevation (ft): 890	Areal water load (m/yr):	0.92
Shore length (m): 2200	Flushing rate $(yr^{-1})$ :	1.30
Watershed area (ha): 14.0	P retention coeff.:	0.90
<pre>% watershed ponded: 0.0</pre>	Lake type: artis	ficial

BIOLOGICAL:	5 January 1993	19 August 1992
DOM. PHYTOPLANKTON (% TOTAL) #3	DINOBRYON 45%	MELOSIRA 60%
#2	SYNEDRA 30%	DINOBRYON 25%
#3		MICROCYSTIS 5%
PHYTOPLANKTON ABUNDANCE (cells/mL)		3765
CHLOROPHYLL-A (µg/L)		12.97
DOM. ZOOPLANKTON (% TOTAL) #3	KERATELLA 63%	KERATELLA 53%
#2	SYNCHAETA 25%	NAUPLIUS LARVA 20%
#3	BOSMINA 12%	COLLOTHECA 10%
ROTIFERS/LITER	115	2670
MICROCRUSTACEA/LITER	16	1199
ZOOPLANKTON ABUNDANCE (#/L)	131	3897
VASCULAR PLANT ABUNDANCE		Sparse
SECCHI DISK TRANSPARENCY (m)		1.0
BOTTOM DISSOLVED OXYGEN (mg/L)	11.3	10.0
BACTERIA (E. coli, #/100 ml) #:		6
#2		4
#:		

#### SUMMER THERMAL STRATIFICATION:

#### not stratified

Depth of thermocline (m): None Hypolimnion volume  $(m^3)$ : None Anoxic volume  $(m^3)$ : None

CHEMICAL:	Lake: COFFIN POND Town: SUGAR HILL				
	5 January 1993		19 August 1992		992
DEPTH (m)	1.0		1.0		
pH (units)	6.6		7.2		
A.N.C. (Alkalinity)	10.6		11.9		
NITRATE NITROGEN	0.10		< 0.02		
TOTAL KJELDAHL NITROGEN	0.24		0.69		
TOTAL PHOSPHORUS	0.012		0.024		
CONDUCTIVITY (µmhos/cm)	150.1		160.5		
APPARENT COLOR (cpu)	13		26		
MAGNESIUM			1.20		
CALCIUM			9.0		
SODIUM			18.3		
POTASSIUM			1.30		
CHLORIDE	34		39		
SULFATE	6		4		
TN : TP	28		29		
CALCITE SATURATION INDEX			2.1		


All results in mg/L unless indicated otherwise

#### TROPHIC CLASSIFICATION: 1992

D.O.	S.D.	PLANT	CHL	TOTAL	CLASS
**	5	0	3	8	Meso.

#### **COMMENTS:**

- 1. Also known as Joe Coffin Lake.
- 2. This is a gravel borrow pit, dug out for the construction of I-93, that filled with water.
- 3. The water appeared turbid with silt; transparency was poor.
- 4. Boat launch was adequate but into very shallow water; parking was good.
- 5. Water level appeared about two feet below normal.
- 6. Bottom of pond was hard gravel with some silt on top.
- 7. The dominant genera of wholewater phytoplankton were the desmids  $\frac{\text{Cosmarium}}{(50\%)}$  and  $\frac{\text{Closterium}}{(25\%)}$ .



### FIELD DATA SHEET

LAKE: COFFIN POND

DATE: 08/19/92

TOWN: SUGAR HILL

	-0	
WEATHER:	OVERCAST,	WARM


DEPTH (M)	TEMP (°C)	*DISSOLVED OXYGEN	OXYGEN SATURATION
0.1	20.0	10.2	112 %
1.0	20.0	10.2	112 %
1.5	20.0	10.0	110 %
	· · · · · · · · · · · · · · · · · · ·		

SECCHI DISK (m): 1.0 COMMENTS:

BOTTOM DEPTH (m): 1.6

TIME: 1125

\*Dissolved oxygen values are in mg/L



## AQUATIC PLANT SURVEY

LAK	E: COFFIN POND	TOWN: SUGAR HILL	DATE: 08/19/92
PLANT		NT NAME	1,51,51,52
Кеу	GENERIC	СОММОИ	ABUNDANCE
Т	Typha	Cattail	Sparse
			,

## OVERALL ABUNDANCE: Sparse

## **GENERAL OBSERVATIONS:**

- 1. Virtually no aquatic plants along the rocky/sandy shoreline.
- 2. Numerous crayfish were observed.